

**HUMAN RPS6KA6-RELATED GENE VARIANT ASSOCIATED  
WITH LUNG CANCERS**

**FIELD OF THE INVENTION**

[0001] The invention relates to the nucleic acid and polypeptide sequences  
5 of a novel human RPS6KA6-related gene variant, the preparation process  
thereof, and the uses of the same in diagnosing cancers, in particular, T cell  
lymphoblastic lymphoma.

**BACKGROUND OF THE INVENTION**

[0002] Lymphoma is the third most common cancer among children in the  
10 world. The major types of lymphoma are Hodgkin's and non-Hodgkin's.  
Non-Hodgkin's lymphoma (NHL) occurs more frequently than Hodgkin's  
disease among children. The major histopathological categories of NHL in  
children are (1) Burkitt's and Burkitt's like lymphomas; (2) lymphoblastic  
lymphomas; (3) anaplastic large cell lymphoma; and (4) diffuse large cell  
15 lymphomas (Percy et al., 1999). In recent years, much progress has been  
made toward understanding the molecular and cellular biology of NHL.  
Many important contributions have been made by the characterization of  
chromosomal translocations and the identification of several key genetic  
factors associated with each type of NHL (Percy et al., 1999). However,  
20 the treatments of NHL still mainly depend on chemotherapy and  
radiotherapy. This is because the molecular mechanisms underlying the  
pathogenesis of NHL remain largely unclear.

[0003] Lymphoblastic lymphoma, a predominant T-cell tumor, accounts  
for about 30% of childhood NHL (National Cancer Institute Cancer.gov  
25 Web site, 2004). Recent studies have shown that T cell lymphoblastic  
lymphoma is caused by abnormal expression of several genetic factors such  
as BCL-6 (Hyjek et al., (2001) Blood. 97: 270-276), MSH2/Lmo-2/Tal-1  
(Lowsky et al., (1997) Blood. 89: 2276-2282) and Stat5 (Kelly et al.,

(2003) J Exp Med. 198: 79-89). Stat5 has been shown to play a role in cell cycle regulation (Nieborowska-Skorska et al., (1999) J Exp Med. 189: 1229-1242; Martino et al., (2001) J Immunol. 166: 1723-1729). Therefore, future strategies for the prevention and treatment of T cell lymphoblastic lymphoma will focus on the elucidation of genetic substrates associated with cell cycle regulation. Interestingly, three members (RSK1, RSK2, and RSK3) of the ribosomal S6 kinase (RSK) family have been shown to be involved in the cell cycle regulation and may play a role in T cell (Edelmann et al., (1996) J Biol Chem. 271: 963-71; Zhao et al., (1996) J Biol Chem. 271: 29773-29779; Brennan et al., (1999) Mol Cell Biol. 19:4729-38; Suzuki et al., (2001) J Immunol. 167:3064-73). Thus, it raised a possibility that RPS6KA6 (ribosomal S6 kinase 4; also named RSK4; GenBank accession # AF184965) has a role in the development of T cell lymphoblastic lymphoma. Therefore, the discovery of gene variants of RPS6KA6 may be important targets for diagnostic markers of T cell lymphoblastic lymphoma.

### **SUMMARY OF THE INVENTION**

[0004] The present invention provides an RPS6KA6-related gene variant (RPS6KA6V) which is negatively expressed in human T cell lymphoblastic lymphoma. The nucleotide sequence of the gene variant and the polypeptide sequence encoded thereby can be used for the diagnosis of any diseases associated with this gene variant or T cell lymphoblastic lymphoma.

[0005] The invention further provides an expression vector and host cell for expressing the variant.

[0006] The invention further provides a method for producing the variant.

[0007] The invention further provides an antibody specifically binding to the variant.

[0008] The invention also provides methods for detecting the presence of the variant in a mammal.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

[0009] FIG. 1 shows the nucleic acid sequence (SEQ ID NO: 1) and amino acid sequence (SEQ ID NO: 2) of RPS6KA6V.

[0010] FIG. 2 shows the nucleotide sequence alignment between the human RPS6KA6 gene (SEQ ID NO: 3) and its related gene variant (RPS6KA6V).

[0011] FIG. 3 shows the amino acid sequence alignment between the human RPS6KA6 protein (SEQ ID NO: 4) and its related gene variant (RPS6KA6V).

[0012] FIG. 4 shows the semi-quantitative RT-PCR analysis of RPS6KA6V in human cell lines, wherein the left and right columns are 100 bp DNA markers.

[0013] FIG. 5 shows the semi-quantitative RT-PCR analysis of RPS6KA6V in human tissue samples, wherein the left and right columns are 100 bp DNA markers.

### **DETAILED DESCRIPTION OF THE INVENTION**

[0014] According to the present invention, all technical and scientific terms used have the same meanings as commonly understood by persons skilled in the art.

[0015] The term "antibody" used herein denotes intact molecules (a polypeptide or group of polypeptides) as well as fragments thereof, such as Fab, R(ab')<sub>2</sub>, and Fv fragments, which are capable of binding the epitopic determinant. Antibodies are produced by specialized B cells after stimulation by an antigen. Structurally, antibody consists of four subunits

including two heavy chains and two light chains. The internal surface shape and charge distribution of the antibody binding domain is complementary to the features of an antigen. Thus, antibody can specifically act against the antigen in an immune response.

5 [0016] The term "base pair (bp)" used herein denotes nucleotides composed of a purine on one strand of DNA which can be hydrogen bonded to a pyrimidine on the other strand. Thymine (or uracil) and adenine residues are linked by two hydrogen bonds. Cytosine and guanine residues are linked by three hydrogen bonds.

10 [0017] The term "Basic Local Alignment Search Tool (BLAST; Altschul et al., (1997) Nucleic Acids Res. 25: 3389-3402)" used herein denotes programs for evaluation of homologies between a query sequence (amino or nucleic acid) and a test sequence as described by Altschul et al. (Nucleic Acids Res. 25: 3389-3402, 1997). Specific BLAST programs are described  
15 as follows:

[0018] (1) BLASTN compares a nucleotide query sequence with a nucleotide sequence database;

[0019] (2) BLASTP compares an amino acid query sequence with a protein sequence database;

20 [0020] (3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence with a protein sequence database;

[0021] (4) TBLASTN compares a query protein sequence with a nucleotide sequence database translated in all six reading frames; and

[0022] (5) TBLASTX compares the six-frame translations of a  
25 nucleotide query sequence with the six-frame translations of a nucleotide sequence database.

[0023] The term "cDNA" used herein denotes nucleic acids that synthesized from a mRNA template using reverse transcriptase.

[0024] The term "cDNA library" used herein denotes a library composed of complementary DNAs, which are reverse-transcribed from mRNAs.

5 [0025] The term "complement" used herein denotes a polynucleotide sequence capable of forming base pairing with another polynucleotide sequence. For example, the sequence 5'-ATGGACTTACT-3' binds to the complementary sequence 5'- AGTAAGTCCAT-3'.

10 [0026] The term "deletion" used herein denotes a removal of a portion of one or more amino acid residues/nucleotides from a gene.

[0027] The term "expressed sequence tags (ESTs)" used herein denotes short (200 to 500 base pairs) nucleotide sequence derived from either 5' or 3' end of a cDNA.

15 [0028] The term "expression vector" used herein denotes nucleic acid constructs which contain a cloning site for introducing the DNA into vector, one or more selectable markers for selecting vectors containing the DNA, an origin of replication for replicating the vector whenever the host cell divides, a terminator sequence, a polyadenylation signal, and a suitable control sequence which can effectively express the DNA in a suitable host.  
20 The suitable control sequence may include promoter, enhancer and other regulatory sequences necessary for directing polymerases to transcribe the DNA.

[0029] The term "host cell" used herein denotes a cell, which is used to receive, maintain, and allow the reproduction of an expression vector comprising DNA. Host cells are transformed or transfected with suitable  
25 vectors constructed using recombinant DNA methods. The recombinant DNA introduced with the vector is replicated whenever the cell divides.

[0030] The term "insertion" or "addition" used herein denotes the addition of a portion of one or more amino acid residues/nucleotides to a gene.

[0031] The term "in silico" used herein denotes a process of using computational methods (e.g., BLAST) to analyze DNA sequences.

5 [0032] The term "polymerase chain reaction (PCR)" used herein denotes a method which increases the copy number of a nucleic acid sequence using a DNA polymerase and a set of primers (about 20bp oligonucleotides complementary to each strand of DNA) under suitable conditions (successive rounds of primer annealing, strand elongation, and  
10 dissociation).

[0033] The term "protein" or "polypeptide" used herein denotes a sequence of amino acids in a specific order that can be encoded by a gene or by a recombinant DNA. It can also be chemically synthesized.

15 [0034] The term "nucleic acid sequence" or "polynucleotide" used herein denotes a sequence of nucleotide (guanine, cytosine, thymine or adenine) in a specific order that can be a natural or synthesized fragment of DNA or RNA. It may be single-stranded or double-stranded.

[0035] The term "reverse transcriptase-polymerase chain reaction (RT-PCR)" used herein denotes a process which transcribes mRNA to  
20 complementary DNA strand using reverse transcriptase followed by polymerase chain reaction to amplify the specific fragment of DNA sequences.

[0036] The term "transformation" used herein denotes a process describing the uptake, incorporation, and expression of exogenous DNA by  
25 prokaryotic host cells.

[0037] The term "transfection" used herein denotes a process describing the uptake, incorporation, and expression of exogenous DNA by eukaryotic host cells.

[0038] The term "variant" used herein denotes a fragment of sequence (nucleotide or amino acid) inserted or deleted by one or more nucleotides/amino acids.

[0039] The present invention provides the polypeptides of a novel human RPS6KA6-related gene variant, as well as the nucleic acid sequences encoding the same.

[0040] According to the present invention, human RPS6KA6 cDNA sequence was used to query the human lung EST databases (a normal lung, a large cell lung cancer, a squamous cell lung cancer and a small cell lung cancer) using BLAST program to search for RPS6KA6-related gene variants. Four ESTs showing similarity to RPS6KA6 were identified. Two were from the large cell lung cancer, one was from the squamous cell lung cancer and one was from the SCLC databases. Their corresponding cDNA clones were found to be identical after sequencing and named RPS6KA6V (RPS6KA6 variant). FIG. 1 shows the nucleic acid sequence of RPS6KA6V (SEQ ID NO: 1) and the amino acid sequence encoded thereby (SEQ ID NO: 2).

[0041] The full-length of the RPS6KA6V cDNA is a 2403bp clone containing a 2094bp open reading frame (ORF) extending from 6bp to 2099bp, which corresponds to an encoded protein of 698 amino acid residues with a predicted molecular mass of 78.2 kDa. To determine the variation in sequence of RPS6KA6V cDNA clone, an alignment of RPS6KA6 nucleotide/amino acid sequence with RPS6KA6V was performed (FIGs. 2 and 3). The results indicate that one major genetic deletion was found in the aligned sequences showing that RPS6KA6V is a 141bp deletion in the sequence of RPS6KA6 from 1978-2118bp. The lack

of 141bp (corresponding to 47aa) is an in-frame deletion in the amino acid sequence of RPS6KA6 and generates a polypeptide of 698 amino acid residues of RPS6KA6V (Fig. 3).

[0042] In the present invention, a search of ESTs deposited in dbEST (Boguski et al. (1993) Nat Genet. 4: 332-3) at NCBI was performed to determine the tissue distribution of RPS6KA6V *in silico*. The result of *in silico* Northern analysis showed that one EST (GenBank accession number AA626690) was found to confirm the absence of 141bp region on RPS6KA6V nucleotide sequence. This EST was also generated from a lung carcinoma cDNA library suggesting that the absence of 141bp nucleotide fragment located between 1977-1978bp of RPS6KA6V may serve as a useful marker for diagnosing cancers associated with this gene variant. Therefore, any nucleotide fragments comprising 1977-1978bp of RPS6KA6V may be used as probes for determining the presence of RPS6KA6V under high stringency conditions. An alternative approach is that any set of primers for amplifying the fragment containing 1977-1978bp of RPS6KA6V may be used for determining the presence of the variant.

[0043] According to the present invention, the polypeptides of the human RPS6KA6V may be produced through genetic engineering techniques. In this case, they are produced by appropriate host cells, which have been transformed by DNAs that code for the polypeptides. The nucleotide sequence encoding the polypeptide containing 657-658aa of the human RPS6KA6V is inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence in a suitable host. The nucleic acid sequence is inserted into the vector in a manner that it will be expressed under appropriate conditions (e.g., in proper orientation and correct reading frame and with appropriate expression sequences, including an RNA polymerase binding sequence and a ribosomal binding sequence).



[0044] Any method that is known to those skilled in the art may be used to construct expression vectors containing sequences encoding the polypeptide of the human RPS6KA6V and appropriate transcriptional/translational control elements. These methods may include *in vitro* recombinant DNA and synthetic techniques, and *in vivo* genetic recombinants. (See, e.g., Sambrook, J. Cold Spring Harbor Press, Plainview N.Y., ch. 4, 8, and 16-17; Ausubel, R. M. et al. (1995) Current protocols in Molecular Biology, John Wiley & Sons, New York N.Y., ch. 9, 13, and 16.)

[0045] A variety of expression vector/host systems may be utilized to express the polypeptide-coding sequence. These include, but not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vector; yeast transformed with yeast expression vector; insect cell systems infected with virus (e.g., baculovirus); plant cell system transformed with viral expression vector (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV); or animal cell system infected with virus (e.g., vaccinia virus, adenovirus, etc.). Preferably, the host cell is a bacterium, and most preferably, the bacterium is *E. coli*.

[0046] Alternatively, the polypeptide of the human RPS6KA6V or the fragments thereof may be synthesized by using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J. Y. et al. (1995) Science 269: 202 to 204). Automated synthesis may be achieved by using the ABI 431A peptide synthesizer (Perkin-Elmer).

[0047] According to the present invention, the polypeptide and nucleic acid sequence of the human RPS6KA6V can be used as immunogen and template of primers/or probes, respectively.

[0048] The present invention further provides the antibodies which specifically bind one or more out-surface epitopes of the polypeptides of the human RPS6KA6V.

[0049] According to the present invention, immunization of mammals with immunogens described herein, preferably humans, rabbits, rats, mice, sheep, goats, cows, or horses, is performed following procedures well known to those skilled in the art, for the purpose of obtaining antisera containing polyclonal antibodies or hybridoma lines secreting monoclonal antibodies.

[0050] Monoclonal antibodies can be prepared by standard techniques, given the teachings contained herein. Such techniques are disclosed, for example, in U.S. Patent Nos. 4,271,145 and 4,196,265. Briefly, an animal is immunized with the immunogen. Hybridomas are prepared by fusing spleen cells from the immunized animal with myeloma cells. The fusion products are screened for those producing antibodies that bind to the immunogen. The positive hybridoma clones are isolated, and the monoclonal antibodies are recovered from those clones.

[0051] Immunization regimens for production of both polyclonal and monoclonal antibodies are well-known in the art. The immunogen may be injected by any of a number of routes, including subcutaneous, intravenous, intraperitoneal, intradermal, intramuscular, mucosal, or a combination thereof. The immunogen may be injected in soluble form, aggregate form, attached to a physical carrier, or mixed with an adjuvant, using methods and materials well-known in the art. The antisera and antibodies may be purified using column chromatography methods well known to those skilled in the art.

[0052] According to the present invention, antibody fragments which contain specific binding sites for the polypeptides may also be generated. For example, such fragments include, but are not limited to, F(ab')<sub>2</sub>

fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments.

[0053] The subject invention also provides methods for diagnosing the diseases associated with the human RPS6KA6V or T cell lymphoblastic lymphoma, by the utilization of the nucleic acid sequence, the polypeptide of the human RPS6KA6V, and the antibodies against the polypeptide.

[0054] Many gene variants have been found to be associated with diseases (Stallings-Mann et al., (1996) Proc Natl Acad Sci U S A 93: 12394-9; Liu et al., (1997) Nat Genet 16:328-9; Siffert et al., (1998) Nat Genet 18: 45 to 8; Lukas et al., (2001) Cancer Res 61: 3212 to 9). Since RPS6KA6V clone was isolated from lung cancers cDNA libraries and identified its expression in lung carcinoma cDNA library using *in silico* Northern analysis, it is advisable that RPS6KA6V may serve as a marker for the diagnosis of human cancers. Thus, the expression level of RPS6KA6V relative to RPS6KA6 may be a useful indicator for screening of patients suspected of having cancers. This suggests that the index of relative expression level (mRNA or protein) may confer an increased susceptibility to cancers. Fragments of RPS6KA6 mRNA may be detected by RT-PCR approach. Polypeptides of RPS6KA6V may be determined by the binding of antibodies to these polypeptides. These approaches may be performed in accordance with conventional methods well known to persons skilled in the art.

[0055] According to the present invention, the expression of the gene variant mRNA in sample may be determined by, but not limited to, RT-PCR. Using TRIZOL reagents (Life Technology), total RNA may be isolated from patient samples. Tissue samples (e.g., biopsy samples) are powdered under liquid nitrogen before homogenization. RNA purity and integrity are assessed by absorbance at 260/280 nm and by agarose gel

electrophoresis. A set of primers can be designed to amplify the expected size of specific PCR fragments of RPS6KA6V. For example, one of the primers may be designed to have a sequence comprising the nucleotides of SEQ ID NO: 1 containing nucleotides 1974 to 1979, and the other may be designed to have a sequence complementary to the nucleotides of SEQ ID NO: 1 at any other locations. Alternatively, one of the primers may be designed to have a sequence complementary to the nucleotides of SEQ ID NO: 1 upstream of nucleotide 1977 and the other may be designed to have a sequence comprising the nucleotides of SEQ ID NO: 1 downstream of nucleotide 1978. In this case, both RPS6KA6 and RPS6KA6V will be amplified. The length of the PCR fragment from RPS6KA6V will be 141bp shorter than that from RPS6KA6. PCR fragments are analyzed on a 1% agarose gel using five microliters (10%) of the amplified products. The intensity of the signals may be determined by using the Molecular Analyst program (version 1.4.1; Bio-Rad). Thus, the index of relative expression levels for each co-amplified PCR product may be calculated based on the intensity of signals.

[0056] The RT-PCR experiment may be performed according to the manufacturer's instructions (Boehringer Mannheim). A 50µl reaction mixture containing 2µl total RNA (0.1µg/µl), 1µl each primer (20 pM), 1µl each dNTP (10 mM), 2.5 µl DTT solution (100 mM), 10 µl 5X RT-PCR buffer, 1µl enzyme mixture, and 28.5 µl sterile distilled water may be subjected to the conditions such as reverse transcription at 60°C for 30 minutes followed by 35 cycles of denaturation at 94°C for 2 minutes, annealing at 60°C for 2 minutes, and extension at 68°C for 2 minutes. The RT-PCR analysis may be repeated twice to ensure reproducibility, for a total of three independent experiments.

[0057] The expression of the gene variant can also be analyzed using Northern Blot hybridization approach. Specific fragment of the RPS6KA6V may be amplified by polymerase chain reaction (PCR) using

primer set designed for RT-PCR. The amplified PCR fragment may be labeled and serve as a probe to hybridize the membranes containing total RNAs extracted from the samples under the conditions of 55°C in a suitable hybridization solution for 3 hr. Blots may be washed twice in 2 x SSC, 0.1% SDS at room temperature for 15 minutes each, followed by two washes in 0.1 x SSC and 0.1% SDS at 65°C for 20 minutes each. After these washes, blot may be rinsed briefly in suitable washing buffer and incubated in blocking solution for 30 minutes, and then incubated in suitable antibody solution for 30 minutes. Blots may be washed in washing buffer for 30 minutes and equilibrated in suitable detection buffer before detecting the signals. Alternatively, the presence of gene variant (cDNAs or PCR) can be detected using microarray approach. The cDNAs or PCR products corresponding to the nucleotide sequences of the present invention may be immobilized on a suitable substrate such as a glass slide. Hybridization can be preformed using the labeled mRNAs extracted from samples. After hybridization, nonhybridized mRNAs are removed. The relative abundance of each labeled transcript, hybridizing to a cDNA/PCR product immobilized on the microarray, can be determined by analyzing the scanned images.

[0058] According to the present invention, the presence of the polypeptide of the gene variant in samples may be determined by, but not limited to, the immunoassay, which uses the antibody specifically binding to the polypeptide. For instance, the polypeptide in protein samples obtained from the mammal suspected of having such diseases may be determined by, but not limited to, the immunoassay wherein the antibody specifically binding to the polypeptide of the invention is brought into contact with the protein samples, and the antibody-polypeptide complex is detected. If necessary, the amount of antibody-polypeptide complex can be determined.

[0059] The polypeptides of the human RPS6KA6V may be expressed in prokaryotic cells by using suitable prokaryotic expression vectors. The

cDNA fragments of RPS6KA6V gene encoding the amino acid coding sequence may be PCR amplified using primer set with restriction enzyme digestion sites incorporated in the 5' and 3' ends, respectively. The PCR products can then be enzyme digested, purified, and inserted into the  
5 corresponding sites of prokaryotic expression vector in-frame to generate recombinant plasmids. Sequence fidelity of this recombinant DNA can be verified by sequencing. The prokaryotic recombinant plasmids may be transformed into host cells (e.g., *E. coli* BL21 (DE3)). Recombinant protein synthesis may be stimulated by the addition of 0.4 mM  
10 isopropylthiogalactoside (IPTG) for 3h. The bacterially-expressed proteins may be purified.

[0060] The polypeptide of the gene variant may be expressed in animal cells by using eukaryotic expression vectors. Cells may be maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10%  
15 fetal bovine serum (FBS; Gibco BRL) at 37°C in a humidified 5% CO<sub>2</sub> atmosphere. Before transfection, the nucleotide sequence of each of the gene variant may be amplified with PCR primers containing restriction enzyme digestion sites and ligated into the corresponding sites of eukaryotic expression vector in-frame. Sequence fidelity of this  
20 recombinant DNA can be verified by sequencing. The cells may be plated in 12-well plates one day before transfection at a density of  $5 \times 10^4$  cells per well. Transfections may be carried out using Lipofectamine Plus transfection reagent according to the manufacturer's instructions (Gibco BRL). Three hours following transfection, medium containing the  
25 complexes may be replaced with fresh medium. Forty-eight hours after incubation, the cells may be scraped into lysis buffer (0.1 M Tris HCl, pH 8.0, 0.1% Triton X-100) for purification of expressed proteins. After these proteins are purified, monoclonal antibodies against these purified proteins (RPS6KA6V) may be generated using hybridoma technique according to  
30 the conventional methods (de StGroth and Scheidegger, (1980) J Immunol

Methods 35:1-21; Cote et al. (1983) Proc Natl Acad Sci U S A 80: 2026-30; and Kozbor et al. (1985) J Immunol Methods 81:31-42).

[0061] According to the present invention, the presence of the polypeptides of the gene variant in samples may be determined by, but not limited to, Western blot analysis. Proteins extracted from samples may be separated by SDS-PAGE and transferred to suitable membranes such as polyvinylidene difluoride (PVDF) in transfer buffer (25 mM Tris-HCl, pH 8.3, 192 mM glycine, 20% methanol) with a Trans-Blot apparatus for 1h at 100 V (e.g., Bio-Rad). The proteins can be immunoblotted with specific antibodies. For example, membrane blotted with extracted proteins may be blocked with suitable buffers such as 3% solution of BSA or 3% solution of nonfat milk powder in TBST buffer (10 mM Tris-HCl, pH 8.0, 150 mM NaCl, 0.1% Tween 20) and incubated with monoclonal antibody directed against the polypeptides of gene variants. Unbound antibody is removed by washing with TBST for 5 X 1 minutes. Bound antibody may be detected using commercial ECL Western blotting detecting reagents.

[0062] The following examples are provided for illustration, but not for limiting the invention.

## **EXAMPLES**

### **Analysis of Human Lung EST Databases**

[0063] Expressed sequence tags (ESTs) generated from the large-scale PCR-based sequencing of the 5'-end of human lung (normal, SCLC, squamous cell lung cancer and large cell lung cancer) cDNA clones were compiled and served as EST databases. Sequence comparisons against the nonredundant nucleotide and protein databases were performed using BLASTN and BLASTX programs (Altschul et al., (1997) Nucleic Acids Res. 25: 3389-3402; Gish and States, (1993) Nat Genet 3:266-272), at the National Center for Biotechnology Information (NCBI) with a significant

cutoff of  $p < 10^{-10}$ . ESTs representing putative RPS6KA6V gene were identified during the course of EST generation.

### **Isolation of cDNA Clones**

[0064] Four identical cDNA clones exhibiting EST sequences similar to the RPS6KA6 gene were isolated from lung cancer cDNA libraries and named RPS6KA6V. The inserts of these clones were subsequently excised *in vivo* from the  $\lambda$ ZAP Express vector using the ExAssist/XLORL helper phage system (Stratagene). Phagemid particles were excised by coinfecting XL1-BLUE MRF' cells with ExAssist helper phage. The excised pBluescript phagemids were used to infect *E. coli* XLORL cells, which lack the amber suppressor necessary for ExAssist phage replication. Infected XLORL cells were selected using kanamycin resistance. Resultant colonies contained the double stranded phagemid vector with the cloned cDNA insert. A single colony was grown overnight in LB-kanamycin, and DNA was purified using a Qiagen plasmid purification kit.

### **Full Length Nucleotide Sequencing and Database Comparisons**

[0065] Phagemid DNA was sequenced using the Epicentre#SE9101LC SequiTherm EXCEL<sup>TM</sup>II DNA Sequencing Kit for 4200S-2 Global NEW IR<sup>2</sup> DNA sequencing system (LI-COR). Using the primer-walking approach, full-length sequence was determined. Nucleotide and protein searches were performed using BLAST against the non-redundant database of NCBI.

### **In Silico Tissue Distribution (Northern) Analysis**

[0066] The coding sequence for each cDNA clones was searched against the dbEST sequence database (Boguski et al., (1993) Nat Genet. 4: 332-3) using the BLAST algorithm at the NCBI website. ESTs derived from each tissue were used as a source of information for transcript tissue expression analysis. Tissue distribution for each isolated cDNA clone was determined



by ESTs matching to that particular sequence variants (insertions or deletions) with a significance cutoff of  $p < 10^{-10}$ .

### **Reverse Transcriptase-Polymerase Chain Reaction (RT-PCR) Analysis**

[0067] Total RNA was extracted from WI-38 (fibroblast), A549 (lung adenocarcinoma), NCI-H661 (lung large cell carcinoma), NCI-H520 (lung squamous carcinoma), NCI-H209 (lung small cell carcinoma), JHH-4 (hepatoma), SUP-T1 (T-cell lymphoblastic lymphoma), Daudi (Burkitt's lymphoma), Ramos (Burkitt's lymphoma), and Raji (Burkitt's lymphoma) cell lines and from four breast cancer, two gastric ulcer, two colon cancer, two hepatoma, one Grave's disease, one colon cancer, one pancreatic carcinoma, one left neck tumor, one gastric carcinoma, two thyroid tumor, one spleen, one pancreatic abscess, one Gastric carcinoma, one adenomatous polyposis, one right neck lymph tissue, one liver cirrhosis, and two parotid gland mixed tumor biopsied samples, RNA purity and integrity were assessed by the absorbance at 260/280 nm and by agarose gel electrophoresis.

[0068] The forward and reverse primers for RPS6KA6V were 5'-GGAGCAAAGGGAGCAATGGTTG-3' (SEQ ID NO: 5) and 5'-TCTTCATCCAGTTTGGCCTAGG-3' (SEQ ID NO: 6), respectively. The expected size of the specific PCR fragment was 170 bp. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH; accession No. M33197) was used as an internal control. The forward and reverse primers for GAPDH were 5'-TGGGTGTGAACCATGAGAAG-3' (SEQ ID NO: 7) and 5'-GTGTCGCTGTTGAAGTCAGA-3' (SEQ ID NO: 8), respectively. The expected size of the PCR fragment was 472 bp. The electrophoresis results of the RPS6KA6V mRNA expression patterns in 10 cell lines and 25 biopsied samples determined by RT-PCR are shown in Figs. 4 and 5. The results showed that RPS6KA6V mRNA was consistently expressed in all cell lines and tissues investigated except in the T-cell lymphoblastic

lymphoma cell line. This suggests that RPS6KA6V can be used for diagnosing T-cell lymphoblastic lymphoma when RPS6KA6V mRNA cannot be detected.

## REFERENCES

Altschul et al., Gapped BLAST and PSI-BLAST: a new generation of protein database search programs, Nucleic Acids Res, 25: 3389-3402, (1997).

Ausubel et al., Current protocols in Molecular Biology, John Wiley & Sons, New York N.Y., ch. 9, 13, and 16, (1995).

Boguski et al., dbEST--database for "expressed sequence tags". Nat Genet. 4: 332-3, (1993).

Brennan et al., p70(s6k) integrates phosphatidylinositol 3-kinase and rapamycin-regulated signals for E2F regulation in T lymphocytes. Mol Cell Biol. 19:4729-38, (1999).

Cote et al., Generation of human monoclonal antibodies reactive with cellular antigens, Proc Natl Acad Sci U S A 80: 2026-30 (1983).

de StGroth and Scheidegger, Production of monoclonal antibodies: strategy and tactics, J Immunol Methods 35:1-21, (1980).

Edelmann et al., Cell cycle regulation of p70 S6 kinase and p42/p44 mitogen-activated protein kinases in Swiss mouse 3T3 fibroblasts. J Biol Chem. 271: 963-71, (1996).

Frodin and Gammeltoft, Role and regulation of 90 kDa ribosomal S6 kinase (RSK) in signal transduction. Mol Cell Endocrinol, 151:65-77, (1999).

Gish and States, Identification of protein coding regions by database similarity search, Nat Genet, 3:266-272, (1993).

Hyjek et al., BCL-6 protein is expressed in precursor T-cell lymphoblastic lymphoma and in prenatal and postnatal thymus. Blood. 97: 270-276 (2001).

Kelly et al., Stat5 synergizes with T cell receptor/antigen stimulation in the development of lymphoblastic lymphoma. *J Exp Med.* 198: 79-89, (2003).

5 Kozbor et al., Specific immunoglobulin production and enhanced tumorigenicity following ascites growth of human hybridomas, *J Immunol Methods*, 81:31-42 (1985).

10 Liu et al., Silent mutation induces exon skipping of fibrillin-1 gene in Marfan syndrome. *Nat Genet* 16:328-9, (1997).

15 Lowsky et al., Defects of the mismatch repair gene MSH2 are implicated in the development of murine and human lymphoblastic lymphomas and are associated with the aberrant expression of rhombotin-2 (Lmo-2) and Tal-1 (SCL). *Blood.* 89: 2276-2282, (1997).

Lukas et al., Alternative and aberrant messenger RNA splicing of the mdm2 oncogene in invasive breast cancer. *Cancer Res* 61:3212-9, (2001).

20 Martino et al., *J Immunol.* Stat5 and Sp1 regulate transcription of the cyclin D2 gene in response to IL-2. 166: 1723-9, (2001).

25 National Cancer Institute Cancer.gov Web site ([http://www.nci.nih.gov/cancerinfo/pdq/treatment/child-nonhodgkins/healthprofessional#Section\\_14](http://www.nci.nih.gov/cancerinfo/pdq/treatment/child-nonhodgkins/healthprofessional#Section_14)). Childhood Non-Hodgkin's Lymphoma (PDQ®): Treatment (Last Modified: 02/18/2004)

30 Nieborowska-Skorska et al., Signal transducer and activator of transcription (STAT)5 activation by BCR/ABL is dependent on intact Src homology (SH)3 and SH2 domains of BCR/ABL and is required for leukemogenesis. *J Exp Med.* 189: 1229-1242, (1999).

35 Percy CL, Smith MA, Linet M, et al.: Lymphomas and reticuloendothelial neoplasms. In: Ries LA, Smith MA, Gurney JG, et al., eds.: *Cancer incidence and survival among children and adolescents: United States SEER Program 1975-1995*. Bethesda, Md: National Cancer Institute, SEER Program, 1999. NIH Pub.No. 99-4649. Also available online. Last accessed

September 03, 2003, pp 35-50. Last accessed April 21, 2003.

Roberge et al., A strategy for a convergent synthesis of N-linked glycopeptides on a solid support. *Science* 269:202-4, (1995).

5

Sambrook, J. Cold Spring Harbor Press, Plainview N.Y., ch. 4, 8, and 16-17.

10

Siffert et al., Association of a human G-protein beta3 subunit variant with hypertension. *Nat Genet*, 18:45-8, (1998).

15

Stallings-Mann et al., Alternative splicing of exon 3 of the human growth hormone receptor is the result of an unusual genetic polymorphism. *Proc Natl Acad Sci U S A* 93:12394-9, (1996).

Suzuki et al., Diverse transcriptional response of CD4(+) T cells to stromal cell-derived factor (SDF)-1: cell survival promotion and priming effects of SDF-1 on CD4(+) T cells. *J Immunol*. 167:3064-73, (2001).

20

Wilson, R.K. GenBank accession # AA626690

Yntema et al., GenBank accession # AF184965

25

Yntema et al., A novel ribosomal S6-kinase (RSK4; RPS6KA6) is commonly deleted in patients with complex X-linked mental retardation. *Genomics*, 62:332-43, (1999).

Zhao et al., Regulation and interaction of pp90(rsk) isoforms with mitogen-activated protein kinases. *J Biol Chem*. 271: 29773-29779, (1996).

30

## SEQUENCE LISTING

&lt;110&gt; Dai, Ken-Shwo

&lt;120&gt; HUMAN RPS6KA6-RELATED GENE VARIANT ASSOCIATED WITH LUNG CANCERS

&lt;130&gt; U 015086-3

&lt;140&gt; --

&lt;141&gt; 2004-03-23

&lt;160&gt; 8

&lt;170&gt; PatentIn version 3.1

&lt;210&gt; 1

&lt;211&gt; 2403

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (6)..(2099)

&lt;223&gt;

&lt;400&gt; 1

```

gggaa atg cta cca ttc gct cct cag gac gag ccc tgg gac cga gaa atg      50
      Met Leu Pro Phe Ala Pro Gln Asp Glu Pro Trp Asp Arg Glu Met
        1             5             10             15

gaa gtg ttc agc ggc ggc ggc gcg agc agc ggc gag gta aat ggt ctt      98
Glu Val Phe Ser Gly Gly Gly Ala Ser Ser Gly Glu Val Asn Gly Leu
              20             25             30

aaa atg gtt gat gag cca atg gaa gag gga gaa gca gat tct tgt cat      146
Lys Met Val Asp Glu Pro Met Glu Glu Gly Glu Ala Asp Ser Cys His
              35             40             45

gat gaa gga gtt gtt aaa gaa atc cct att act cat cat gtt aag gaa      194
Asp Glu Gly Val Val Lys Glu Ile Pro Ile Thr His His Val Lys Glu
              50             55             60

ggc tat gag aaa gca gat cct gca cag ttt gag ttg ctc aag gtt ctt      242
Gly Tyr Glu Lys Ala Asp Pro Ala Gln Phe Glu Leu Leu Lys Val Leu
              65             70             75

ggt cag ggg tca ttt gga aag gtt ttt ctt gtt aga aag aag acc ggt      290
Gly Gln Gly Ser Phe Gly Lys Val Phe Leu Val Arg Lys Lys Thr Gly
              80             85             90             95

cct gat gct ggg cag ctc tat gca atg aag gtg tta aaa aaa gcc tct      338
Pro Asp Ala Gly Gln Leu Tyr Ala Met Lys Val Leu Lys Lys Ala Ser
              100            105            110

tta aaa gtt cga gac aga gtt cgg aca aag atg gag agg gat ata ctg      386
Leu Lys Val Arg Asp Arg Val Arg Thr Lys Met Glu Arg Asp Ile Leu
              115            120            125

gtg gaa gta aat cat cca ttt att gtc aaa ttg cac tat gcc ttt cag      434
Val Glu Val Asn His Pro Phe Ile Val Lys Leu His Tyr Ala Phe Gln
              130            135            140

```

act gaa ggg aaa ctg tac tta ata ctg gat ttt ctc agg gga gga gat	482
Thr Glu Gly Lys Leu Tyr Leu Ile Leu Asp Phe Leu Arg Gly Gly Asp	
145 150 155	
ggt ttc aca aga tta tcc aaa gag gtt ctg ttt aca gag gaa gat gtg	530
Val Phe Thr Arg Leu Ser Lys Glu Val Leu Phe Thr Glu Glu Asp Val	
160 165 170 175	
aaa ttc tac ctc gca gaa ctg gcc ctt gct ttg gat cat ctg cac caa	578
Lys Phe Tyr Leu Ala Glu Leu Ala Leu Ala Leu Asp His Leu His Gln	
180 185 190	
tta gga att gtt tat aga gac ctg aag cca gaa aac att ttg ctt gat	626
Leu Gly Ile Val Tyr Arg Asp Leu Lys Pro Glu Asn Ile Leu Leu Asp	
195 200 205	
gaa ata gga cat atc aaa tta aca gat ttt gga ctc agc aag gag tca	674
Glu Ile Gly His Ile Lys Leu Thr Asp Phe Gly Leu Ser Lys Glu Ser	
210 215 220	
gta gat caa gaa aag aag gct tac tca ttt tgt ggt aca gta gag tat	722
Val Asp Gln Glu Lys Lys Ala Tyr Ser Phe Cys Gly Thr Val Glu Tyr	
225 230 235	
atg gct cct gaa gta gta aat agg aga ggc cat tcc cag agt gct gat	770
Met Ala Pro Glu Val Val Asn Arg Arg Gly His Ser Gln Ser Ala Asp	
240 245 250 255	
tgg tgg tca tat ggt gtt ctt atg ttt gaa atg ctt act ggt act ctg	818
Trp Trp Ser Tyr Gly Val Leu Met Phe Glu Met Leu Thr Gly Thr Leu	
260 265 270	
cca ttt caa ggt aaa gac aga aat gag acc atg aat atg ata tta aaa	866
Pro Phe Gln Gly Lys Asp Arg Asn Glu Thr Met Asn Met Ile Leu Lys	
275 280 285	
gca aaa ctt gga atg cct caa ttt ctt agt gct gaa gca caa agt ctt	914
Ala Lys Leu Gly Met Pro Gln Phe Leu Ser Ala Glu Ala Gln Ser Leu	
290 295 300	
cta agg atg tta ttc aaa agg aat cca gca aat aga ttg gga tca gaa	962
Leu Arg Met Leu Phe Lys Arg Asn Pro Ala Asn Arg Leu Gly Ser Glu	
305 310 315	
gga gtt gaa gaa atc aaa aga cat ctg ttt ttt gca aat att gac tgg	1010
Gly Val Glu Glu Ile Lys Arg His Leu Phe Phe Ala Asn Ile Asp Trp	
320 325 330 335	
gat aaa tta tat aaa aga gaa gtt caa cct cct ttc aaa cct gct tct	1058
Asp Lys Leu Tyr Lys Arg Glu Val Gln Pro Pro Phe Lys Pro Ala Ser	
340 345 350	
gga aaa cca gat gat act ttt tgt ttt gat cct gaa ttt act gca aaa	1106
Gly Lys Pro Asp Asp Thr Phe Cys Phe Asp Pro Glu Phe Thr Ala Lys	
355 360 365	
aca cct aaa gat tct ccc ggt ttg cca gcc agt gca aat gct cat cag	1154
Thr Pro Lys Asp Ser Pro Gly Leu Pro Ala Ser Ala Asn Ala His Gln	
370 375 380	

ctc ttc aaa gga ttc agc ttt gtt gca act tct att gca gaa gaa tat	1202
Leu Phe Lys Gly Phe Ser Phe Val Ala Thr Ser Ile Ala Glu Glu Tyr	
385 390 395	
aaa atc act cct atc aca agt gca aat gta tta cca att gtt cag ata	1250
Lys Ile Thr Pro Ile Thr Ser Ala Asn Val Leu Pro Ile Val Gln Ile	
400 405 410 415	
aat gga aat gct gca caa ttt ggt gaa gta tat gaa ttg aag gag gat	1298
Asn Gly Asn Ala Ala Gln Phe Gly Glu Val Tyr Glu Leu Lys Glu Asp	
420 425 430	
att ggt gtt ggc tcc tac tct gtt tgc aag cga tgc ata cat gca act	1346
Ile Gly Val Gly Ser Tyr Ser Val Cys Lys Arg Cys Ile His Ala Thr	
435 440 445	
acc aac atg gaa ttt gca gtg aag atc att gac aaa agt aag cga gac	1394
Thr Asn Met Glu Phe Ala Val Lys Ile Ile Asp Lys Ser Lys Arg Asp	
450 455 460	
cct tca gaa gag att gaa ata ttg atg cgc tat gga caa cat ccc aac	1442
Pro Ser Glu Glu Ile Glu Ile Leu Met Arg Tyr Gly Gln His Pro Asn	
465 470 475	
att att act ttg aag gat gtc ttt gat gat ggt aga tat gtt tac ctt	1490
Ile Ile Thr Leu Lys Asp Val Phe Asp Asp Gly Arg Tyr Val Tyr Leu	
480 485 490 495	
gtt acg gat tta atg aaa gga gga gag tta ctt gac cgt att ctc aaa	1538
Val Thr Asp Leu Met Lys Gly Gly Glu Leu Leu Asp Arg Ile Leu Lys	
500 505 510	
caa aaa tgt ttc tcg gaa cgg gag gct agt gat ata cta tat gta ata	1586
Gln Lys Cys Phe Ser Glu Arg Glu Ala Ser Asp Ile Leu Tyr Val Ile	
515 520 525	
agt aag aca gtt gac tat ctt cat tgt caa gga gtt gtt cat cgt gat	1634
Ser Lys Thr Val Asp Tyr Leu His Cys Gln Gly Val Val His Arg Asp	
530 535 540	
ctt aaa cct agt aat att tta tac atg gat gaa tca gcc agt gca gat	1682
Leu Lys Pro Ser Asn Ile Leu Tyr Met Asp Glu Ser Ala Ser Ala Asp	
545 550 555	
tca atc agg ata tgt gat ttt ggg ttt gca aaa caa ctt cga gga gaa	1730
Ser Ile Arg Ile Cys Asp Phe Gly Phe Ala Lys Gln Leu Arg Gly Glu	
560 565 570 575	
aat gga ctt ctc tta act cca tgc tac act gca aac ttt gtt gca cct	1778
Asn Gly Leu Leu Leu Thr Pro Cys Tyr Thr Ala Asn Phe Val Ala Pro	
580 585 590	
gag gtt ctt atg caa cag gga tat gat gct gct tgt gat atc tgg agt	1826
Glu Val Leu Met Gln Gln Gly Tyr Asp Ala Ala Cys Asp Ile Trp Ser	
595 600 605	
tta gga gtc ctt ttt tac aca atg ttg gct ggc tac act cca ttt gct	1874
Leu Gly Val Leu Phe Tyr Thr Met Leu Ala Gly Tyr Thr Pro Phe Ala	
610 615 620	
aat ggc ccc aat gat act cct gaa gag ata ctg ctg cgt ata ggc aat	1922



Asn	Gly	Pro	Asn	Asp	Thr	Pro	Glu	Glu	Ile	Leu	Leu	Arg	Ile	Gly	Asn		
625						630					635						
gga	aaa	ttc	tct	ttg	agt	ggg	gga	aac	tgg	gac	aat	att	tca	gac	gga	1970	
Gly	Lys	Phe	Ser	Leu	Ser	Gly	Gly	Asn	Trp	Asp	Asn	Ile	Ser	Asp	Gly		
640					645					650					655		
gca	aag	gga	gca	atg	gtt	gca	aca	tac	tct	gcc	ctg	act	cac	aag	acc	2018	
Ala	Lys	Gly	Ala	Met	Val	Ala	Thr	Tyr	Ser	Ala	Leu	Thr	His	Lys	Thr		
				660					665					670			
ttt	caa	cca	gtc	cta	gag	cct	gta	gct	gct	tca	agc	tta	gcc	cag	cga	2066	
Phe	Gln	Pro	Val	Leu	Glu	Pro	Val	Ala	Ala	Ser	Ser	Leu	Ala	Gln	Arg		
			675					680					685				
cgg	agc	atg	aaa	aag	cga	aca	tca	act	ggc	ctg	taagattt	ggtgtt	ccta			2119	
Arg	Ser	Met	Lys	Lys	Arg	Thr	Ser	Thr	Gly	Leu							
		690					695										
ggccaaactg	gatgaagatg	aaattaaatg	tgtggctttt	ttcctattct	tatcaaaggc											2179	
atcgttgtct	gctaaattac	ttgaatatta	agtaatatta	aatccccatt	tttaggggaa											2239	
gtgagattta	aaaaaccatt	cacaggtcca	caatattcat	actatgtgtt	tgtagtagtg											2299	
ttcaagtgtt	tatttaagca	tataattggt	gtccaccagg	tcctcacaac	ttctctgcac											2359	
acaagcttct	aaaattcctt	tcaaataaag	ttactttaat	attt												2403	

  

<210> 2  
 <211> 698  
 <212> PRT  
 <213> Homo sapiens

<400> 2

Met	Leu	Pro	Phe	Ala	Pro	Gln	Asp	Glu	Pro	Trp	Asp	Arg	Glu	Met	Glu		
1				5					10					15			
Val	Phe	Ser	Gly	Gly	Gly	Ala	Ser	Ser	Gly	Glu	Val	Asn	Gly	Leu	Lys		
			20					25					30				
Met	Val	Asp	Glu	Pro	Met	Glu	Glu	Gly	Glu	Ala	Asp	Ser	Cys	His	Asp		
		35					40					45					
Glu	Gly	Val	Val	Lys	Glu	Ile	Pro	Ile	Thr	His	His	Val	Lys	Glu	Gly		
	50					55					60						
Tyr	Glu	Lys	Ala	Asp	Pro	Ala	Gln	Phe	Glu	Leu	Leu	Lys	Val	Leu	Gly		
65					70					75					80		
Gln	Gly	Ser	Phe	Gly	Lys	Val	Phe	Leu	Val	Arg	Lys	Lys	Thr	Gly	Pro		
				85					90					95			

Asp Ala Gly Gln Leu Tyr Ala Met Lys Val Leu Lys Lys Ala Ser Leu  
100 105 110

Lys Val Arg Asp Arg Val Arg Thr Lys Met Glu Arg Asp Ile Leu Val  
115 120 125

Glu Val Asn His Pro Phe Ile Val Lys Leu His Tyr Ala Phe Gln Thr  
130 135 140

Glu Gly Lys Leu Tyr Leu Ile Leu Asp Phe Leu Arg Gly Gly Asp Val  
145 150 155 160

Phe Thr Arg Leu Ser Lys Glu Val Leu Phe Thr Glu Glu Asp Val Lys  
165 170 175

Phe Tyr Leu Ala Glu Leu Ala Leu Ala Leu Asp His Leu His Gln Leu  
180 185 190

Gly Ile Val Tyr Arg Asp Leu Lys Pro Glu Asn Ile Leu Leu Asp Glu  
195 200 205

Ile Gly His Ile Lys Leu Thr Asp Phe Gly Leu Ser Lys Glu Ser Val  
210 215 220

Asp Gln Glu Lys Lys Ala Tyr Ser Phe Cys Gly Thr Val Glu Tyr Met  
225 230 235 240

Ala Pro Glu Val Val Asn Arg Arg Gly His Ser Gln Ser Ala Asp Trp  
245 250 255

Trp Ser Tyr Gly Val Leu Met Phe Glu Met Leu Thr Gly Thr Leu Pro  
260 265 270

Phe Gln Gly Lys Asp Arg Asn Glu Thr Met Asn Met Ile Leu Lys Ala  
275 280 285

Lys Leu Gly Met Pro Gln Phe Leu Ser Ala Glu Ala Gln Ser Leu Leu  
290 295 300

Arg Met Leu Phe Lys Arg Asn Pro Ala Asn Arg Leu Gly Ser Glu Gly  
305 310 315 320

Val Glu Glu Ile Lys Arg His Leu Phe Phe Ala Asn Ile Asp Trp Asp  
325 330 335

Lys Leu Tyr Lys Arg Glu Val Gln Pro Pro Phe Lys Pro Ala Ser Gly

				340						345						350	
Lys	Pro	Asp	Asp	Thr	Phe	Cys	Phe	Asp	Pro	Glu	Phe	Thr	Ala	Lys	Thr		
		355					360					365					
Pro	Lys	Asp	Ser	Pro	Gly	Leu	Pro	Ala	Ser	Ala	Asn	Ala	His	Gln	Leu		
	370					375					380						
Phe	Lys	Gly	Phe	Ser	Phe	Val	Ala	Thr	Ser	Ile	Ala	Glu	Glu	Tyr	Lys		
385					390					395					400		
Ile	Thr	Pro	Ile	Thr	Ser	Ala	Asn	Val	Leu	Pro	Ile	Val	Gln	Ile	Asn		
				405					410					415			
Gly	Asn	Ala	Ala	Gln	Phe	Gly	Glu	Val	Tyr	Glu	Leu	Lys	Glu	Asp	Ile		
			420					425					430				
Gly	Val	Gly	Ser	Tyr	Ser	Val	Cys	Lys	Arg	Cys	Ile	His	Ala	Thr	Thr		
		435					440					445					
Asn	Met	Glu	Phe	Ala	Val	Lys	Ile	Ile	Asp	Lys	Ser	Lys	Arg	Asp	Pro		
	450					455					460						
Ser	Glu	Glu	Ile	Glu	Ile	Leu	Met	Arg	Tyr	Gly	Gln	His	Pro	Asn	Ile		
465					470					475					480		
Ile	Thr	Leu	Lys	Asp	Val	Phe	Asp	Asp	Gly	Arg	Tyr	Val	Tyr	Leu	Val		
				485					490					495			
Thr	Asp	Leu	Met	Lys	Gly	Gly	Glu	Leu	Leu	Asp	Arg	Ile	Leu	Lys	Gln		
			500					505					510				
Lys	Cys	Phe	Ser	Glu	Arg	Glu	Ala	Ser	Asp	Ile	Leu	Tyr	Val	Ile	Ser		
		515					520					525					
Lys	Thr	Val	Asp	Tyr	Leu	His	Cys	Gln	Gly	Val	Val	His	Arg	Asp	Leu		
	530					535					540						
Lys	Pro	Ser	Asn	Ile	Leu	Tyr	Met	Asp	Glu	Ser	Ala	Ser	Ala	Asp	Ser		
545					550					555					560		
Ile	Arg	Ile	Cys	Asp	Phe	Gly	Phe	Ala	Lys	Gln	Leu	Arg	Gly	Glu	Asn		
				565					570					575			
Gly	Leu	Leu	Leu	Thr	Pro	Cys	Tyr	Thr	Ala	Asn	Phe	Val	Ala	Pro	Glu		
			580					585					590				

Val Leu Met Gln Gln Gly Tyr Asp Ala Ala Cys Asp Ile Trp Ser Leu  
595 600 605

Gly Val Leu Phe Tyr Thr Met Leu Ala Gly Tyr Thr Pro Phe Ala Asn  
610 615 620

Gly Pro Asn Asp Thr Pro Glu Glu Ile Leu Leu Arg Ile Gly Asn Gly  
625 630 635 640

Lys Phe Ser Leu Ser Gly Gly Asn Trp Asp Asn Ile Ser Asp Gly Ala  
645 650 655

Lys Gly Ala Met Val Ala Thr Tyr Ser Ala Leu Thr His Lys Thr Phe  
660 665 670

Gln Pro Val Leu Glu Pro Val Ala Ala Ser Ser Leu Ala Gln Arg Arg  
675 680 685

Ser Met Lys Lys Arg Thr Ser Thr Gly Leu  
690 695

<210> 3  
<211> 2235  
<212> DNA  
<213> Homo sapiens

<220>  
<221> CDS  
<222> (1) .. (2235)  
<223>

<400> 3  
atg cta cca ttc gct cct cag gac gag ccc tgg gac cga gaa atg gaa 48  
Met Leu Pro Phe Ala Pro Gln Asp Glu Pro Trp Asp Arg Glu Met Glu  
1 5 10 15  
gtg ttc agc ggc ggc ggc gcg agc agc ggc gag gta aat ggt ctt aaa 96  
Val Phe Ser Gly Gly Gly Ala Ser Ser Gly Glu Val Asn Gly Leu Lys  
20 25 30  
atg gtt gat gag cca atg gaa gag gga gaa gca gat tct tgt cat gat 144  
Met Val Asp Glu Pro Met Glu Glu Gly Glu Ala Asp Ser Cys His Asp  
35 40 45  
gaa gga gtt gtt aaa gaa atc cct att act cat cat gtt aag gaa ggc 192  
Glu Gly Val Val Lys Glu Ile Pro Ile Thr His His Val Lys Glu Gly  
50 55 60  
tat gag aaa gca gat cct gca cag ttt gag ttg ctc aag gtt ctt ggt 240  
Tyr Glu Lys Ala Asp Pro Ala Gln Phe Glu Leu Leu Lys Val Leu Gly  
65 70 75 80

cag ggg tca ttt gga aag gtt ttt ctt gtt aga aag aag acc ggt cct	288
Gln Gly Ser Phe Gly Lys Val Phe Leu Val Arg Lys Lys Thr Gly Pro	
85 90 95	
gat gct ggg cag ctc tat gca atg aag gtg tta aaa aaa gcc tct tta	336
Asp Ala Gly Gln Leu Tyr Ala Met Lys Val Leu Lys Lys Ala Ser Leu	
100 105 110	
aaa gtt cga gac aga gtt cgg aca aag atg gag agg gat ata ctg gtg	384
Lys Val Arg Asp Arg Val Arg Thr Lys Met Glu Arg Asp Ile Leu Val	
115 120 125	
gaa gta aat cat cca ttt att gtc aaa ttg cac tat gcc ttt cag act	432
Glu Val Asn His Pro Phe Ile Val Lys Leu His Tyr Ala Phe Gln Thr	
130 135 140	
gaa ggg aaa ctg tac tta ata ctg gat ttt ctc agg gga gga gat gtt	480
Glu Gly Lys Leu Tyr Leu Ile Leu Asp Phe Leu Arg Gly Gly Asp Val	
145 150 155 160	
ttc aca aga tta tcc aaa gag gtt ctg ttt aca gag gaa gat gtg aaa	528
Phe Thr Arg Leu Ser Lys Glu Val Leu Phe Thr Glu Glu Asp Val Lys	
165 170 175	
ttc tac ctc gca gaa ctg gcc ctt gct ttg gat cat ctg cac caa tta	576
Phe Tyr Leu Ala Glu Leu Ala Leu Ala Leu Asp His Leu His Gln Leu	
180 185 190	
gga att gtt tat aga gac ctg aag cca gaa aac att ttg ctt gat gaa	624
Gly Ile Val Tyr Arg Asp Leu Lys Pro Glu Asn Ile Leu Leu Asp Glu	
195 200 205	
ata gga cat atc aaa tta aca gat ttt gga ctc agc aag gag tca gta	672
Ile Gly His Ile Lys Leu Thr Asp Phe Gly Leu Ser Lys Glu Ser Val	
210 215 220	
gat caa gaa aag aag gct tac tca ttt tgt ggt aca gta gag tat atg	720
Asp Gln Glu Lys Lys Ala Tyr Ser Phe Cys Gly Thr Val Glu Tyr Met	
225 230 235 240	
gct cct gaa gta gta aat agg aga ggc cat tcc cag agt gct gat tgg	768
Ala Pro Glu Val Val Asn Arg Arg Gly His Ser Gln Ser Ala Asp Trp	
245 250 255	
tgg tca tat ggt gtt ctt atg ttt gaa atg ctt act ggt act ctg cca	816
Trp Ser Tyr Gly Val Leu Met Phe Glu Met Leu Thr Gly Thr Leu Pro	
260 265 270	
ttt caa ggt aaa gac aga aat gag acc atg aat atg ata tta aaa gca	864
Phe Gln Gly Lys Asp Arg Asn Glu Thr Met Asn Met Ile Leu Lys Ala	
275 280 285	
aaa ctt gga atg cct caa ttt ctt agt gct gaa gca caa agt ctt cta	912
Lys Leu Gly Met Pro Gln Phe Leu Ser Ala Glu Ala Gln Ser Leu Leu	
290 295 300	
agg atg tta ttc aaa agg aat cca gca aat aga ttg gga tca gaa gga	960
Arg Met Leu Phe Lys Arg Asn Pro Ala Asn Arg Leu Gly Ser Glu Gly	
305 310 315 320	
gtt gaa gaa atc aaa aga cat ctg ttt ttt gca aat att gac tgg gat	1008

Val	Glu	Glu	Ile	Lys	Arg	His	Leu	Phe	Phe	Ala	Asn	Ile	Asp	Trp	Asp	
				325					330					335		
aaa	tta	tat	aaa	aga	gaa	gtt	caa	cct	cct	ttc	aaa	cct	gct	tct	gga	1056
Lys	Leu	Tyr	Lys	Arg	Glu	Val	Gln	Pro	Pro	Phe	Lys	Pro	Ala	Ser	Gly	
			340					345					350			
aaa	cca	gat	gat	act	ttt	tgt	ttt	gat	cct	gaa	ttt	act	gca	aaa	aca	1104
Lys	Pro	Asp	Asp	Thr	Phe	Cys	Phe	Asp	Pro	Glu	Phe	Thr	Ala	Lys	Thr	
		355					360					365				
cct	aaa	gat	tct	ccc	ggg	ttg	cca	gcc	agt	gca	aat	gct	cat	cag	ctc	1152
Pro	Lys	Asp	Ser	Pro	Gly	Leu	Pro	Ala	Ser	Ala	Asn	Ala	His	Gln	Leu	
	370					375					380					
ttc	aaa	gga	ttc	agc	ttt	gtt	gca	act	tct	att	gca	gaa	gaa	tat	aaa	1200
Phe	Lys	Gly	Phe	Ser	Phe	Val	Ala	Thr	Ser	Ile	Ala	Glu	Glu	Tyr	Lys	
385					390					395					400	
atc	act	cct	atc	aca	agt	gca	aat	gta	tta	cca	att	gtt	cag	ata	aat	1248
Ile	Thr	Pro	Ile	Thr	Ser	Ala	Asn	Val	Leu	Pro	Ile	Val	Gln	Ile	Asn	
				405					410					415		
gga	aat	gct	gca	caa	ttt	ggg	gaa	gta	tat	gaa	ttg	aag	gag	gat	att	1296
Gly	Asn	Ala	Ala	Gln	Phe	Gly	Glu	Val	Tyr	Glu	Leu	Lys	Glu	Asp	Ile	
			420					425					430			
ggg	gtt	ggc	tcc	tac	tct	gtt	tgc	aag	cga	tgc	ata	cat	gca	act	acc	1344
Gly	Val	Gly	Ser	Tyr	Ser	Val	Cys	Lys	Arg	Cys	Ile	His	Ala	Thr	Thr	
		435					440					445				
aac	atg	gaa	ttt	gca	gtg	aag	atc	att	gac	aaa	agt	aag	cga	gac	cct	1392
Asn	Met	Glu	Phe	Ala	Val	Lys	Ile	Ile	Asp	Lys	Ser	Lys	Arg	Asp	Pro	
	450					455					460					
tca	gaa	gag	att	gaa	ata	ttg	atg	cgc	tat	gga	caa	cat	ccc	aac	att	1440
Ser	Glu	Glu	Ile	Glu	Ile	Leu	Met	Arg	Tyr	Gly	Gln	His	Pro	Asn	Ile	
465					470					475					480	
att	act	ttg	aag	gat	gtc	ttt	gat	gat	ggg	aga	tat	gtt	tac	ctt	gtt	1488
Ile	Thr	Leu	Lys	Asp	Val	Phe	Asp	Asp	Gly	Arg	Tyr	Val	Tyr	Leu	Val	
				485					490					495		
acg	gat	tta	atg	aaa	gga	gga	gag	tta	ctt	gac	cgt	att	ctc	aaa	caa	1536
Thr	Asp	Leu	Met	Lys	Gly	Gly	Glu	Leu	Leu	Asp	Arg	Ile	Leu	Lys	Gln	
			500					505					510			
aaa	tgt	ttc	tcg	gaa	cgg	gag	gct	agt	gat	ata	cta	tat	gta	ata	agt	1584
Lys	Cys	Phe	Ser	Glu	Arg	Glu	Ala	Ser	Asp	Ile	Leu	Tyr	Val	Ile	Ser	
		515					520					525				
aag	aca	gtt	gac	tat	ctt	cat	tgt	caa	gga	gtt	gtt	cat	cgt	gat	ctt	1632
Lys	Thr	Val	Asp	Tyr	Leu	His	Cys	Gln	Gly	Val	Val	His	Arg	Asp	Leu	
		530				535					540					
aaa	cct	agt	aat	att	tta	tac	atg	gat	gaa	tca	gcc	agt	gca	gat	tca	1680
Lys	Pro	Ser	Asn	Ile	Leu	Tyr	Met	Asp	Glu	Ser	Ala	Ser	Ala	Asp	Ser	
545					550					555					560	
atc	agg	ata	tgt	gat	ttt	ggg	ttt	gca	aaa	caa	ctt	cga	gga	gaa	aat	1728
Ile	Arg	Ile	Cys	Asp	Phe	Gly	Phe	Ala	Lys	Gln	Leu	Arg	Gly	Glu	Asn	

565										570					575					
gga	ctt	ctc	tta	act	cca	tgc	tac	act	gca	aac	ttt	gtt	gca	cct	gag	1776				
Gly	Leu	Leu	Leu	Thr	Pro	Cys	Tyr	Thr	Ala	Asn	Phe	Val	Ala	Pro	Glu					
			580					585					590							
gtt	ctt	atg	caa	cag	gga	tat	gat	gct	gct	tgt	gat	atc	tgg	agt	tta	1824				
Val	Leu	Met	Gln	Gln	Gly	Tyr	Asp	Ala	Ala	Cys	Asp	Ile	Trp	Ser	Leu					
		595					600					605								
gga	gtc	ctt	ttt	tac	aca	atg	ttg	gct	ggc	tac	act	cca	ttt	gct	aat	1872				
Gly	Val	Leu	Phe	Tyr	Thr	Met	Leu	Ala	Gly	Tyr	Thr	Pro	Phe	Ala	Asn					
	610					615					620									
ggc	ccc	aat	gat	act	cct	gaa	gag	ata	ctg	ctg	cgt	ata	ggc	aat	gga	1920				
Gly	Pro	Asn	Asp	Thr	Pro	Glu	Glu	Ile	Leu	Leu	Arg	Ile	Gly	Asn	Gly					
	625					630				635					640					
aaa	ttc	tct	ttg	agt	ggg	gga	aac	tgg	gac	aat	att	tca	gac	gga	gca	1968				
Lys	Phe	Ser	Leu	Ser	Gly	Gly	Asn	Trp	Asp	Asn	Ile	Ser	Asp	Gly	Ala					
				645					650					655						
aag	gat	ttg	ctt	tcc	cat	atg	ctt	cat	atg	gac	cca	cat	cag	cgg	tat	2016				
Lys	Asp	Leu	Leu	Ser	His	Met	Leu	His	Met	Asp	Pro	His	Gln	Arg	Tyr					
			660					665					670							
act	gct	gaa	caa	ata	tta	aag	cac	tca	tgg	ata	act	cac	aga	gac	cag	2064				
Thr	Ala	Glu	Gln	Ile	Leu	Lys	His	Ser	Trp	Ile	Thr	His	Arg	Asp	Gln					
		675					680					685								
ttg	cca	aat	gat	cag	cca	aag	aga	aat	gat	gtg	tca	cat	gtt	gtt	aag	2112				
Leu	Pro	Asn	Asp	Gln	Pro	Lys	Arg	Asn	Asp	Val	Ser	His	Val	Val	Lys					
	690					695					700									
gga	gca	atg	gtt	gca	aca	tac	tct	gcc	ctg	act	cac	aag	acc	ttt	caa	2160				
Gly	Ala	Met	Val	Ala	Thr	Tyr	Ser	Ala	Leu	Thr	His	Lys	Thr	Phe	Gln					
	705				710					715					720					
cca	gtc	cta	gag	cct	gta	gct	gct	tca	agc	tta	gcc	cag	cga	cgg	agc	2208				
Pro	Val	Leu	Glu	Pro	Val	Ala	Ala	Ser	Ser	Leu	Ala	Gln	Arg	Arg	Ser					
				725					730					735						
atg	aaa	aag	cga	aca	tca	act	ggc	ctg								2235				
Met	Lys	Lys	Arg	Thr	Ser	Thr	Gly	Leu												
			740					745												

<210> 4  
 <211> 745  
 <212> PRT  
 <213> Homo sapiens

<400> 4

Met	Leu	Pro	Phe	Ala	Pro	Gln	Asp	Glu	Pro	Trp	Asp	Arg	Glu	Met	Glu
1				5					10					15	

Val	Phe	Ser	Gly	Gly	Gly	Ala	Ser	Ser	Gly	Glu	Val	Asn	Gly	Leu	Lys
			20					25					30		

Met	Val	Asp	Glu	Pro	Met	Glu	Glu	Gly	Glu	Ala	Asp	Ser	Cys	His	Asp		
		35					40					45					
Glu	Gly	Val	Val	Lys	Glu	Ile	Pro	Ile	Thr	His	His	Val	Lys	Glu	Gly		
	50					55					60						
Tyr	Glu	Lys	Ala	Asp	Pro	Ala	Gln	Phe	Glu	Leu	Leu	Lys	Val	Leu	Gly		
65					70					75					80		
Gln	Gly	Ser	Phe	Gly	Lys	Val	Phe	Leu	Val	Arg	Lys	Lys	Thr	Gly	Pro		
				85					90					95			
Asp	Ala	Gly	Gln	Leu	Tyr	Ala	Met	Lys	Val	Leu	Lys	Lys	Ala	Ser	Leu		
			100					105					110				
Lys	Val	Arg	Asp	Arg	Val	Arg	Thr	Lys	Met	Glu	Arg	Asp	Ile	Leu	Val		
		115					120					125					
Glu	Val	Asn	His	Pro	Phe	Ile	Val	Lys	Leu	His	Tyr	Ala	Phe	Gln	Thr		
	130					135					140						
Glu	Gly	Lys	Leu	Tyr	Leu	Ile	Leu	Asp	Phe	Leu	Arg	Gly	Gly	Asp	Val		
145					150					155					160		
Phe	Thr	Arg	Leu	Ser	Lys	Glu	Val	Leu	Phe	Thr	Glu	Glu	Asp	Val	Lys		
			165						170					175			
Phe	Tyr	Leu	Ala	Glu	Leu	Ala	Leu	Ala	Leu	Asp	His	Leu	His	Gln	Leu		
		180						185					190				
Gly	Ile	Val	Tyr	Arg	Asp	Leu	Lys	Pro	Glu	Asn	Ile	Leu	Leu	Asp	Glu		
		195					200					205					
Ile	Gly	His	Ile	Lys	Leu	Thr	Asp	Phe	Gly	Leu	Ser	Lys	Glu	Ser	Val		
	210					215					220						
Asp	Gln	Glu	Lys	Lys	Ala	Tyr	Ser	Phe	Cys	Gly	Thr	Val	Glu	Tyr	Met		
225					230					235					240		
Ala	Pro	Glu	Val	Val	Asn	Arg	Arg	Gly	His	Ser	Gln	Ser	Ala	Asp	Trp		
			245						250					255			
Trp	Ser	Tyr	Gly	Val	Leu	Met	Phe	Glu	Met	Leu	Thr	Gly	Thr	Leu	Pro		
			260					265					270				



Phe Gln Gly Lys Asp Arg Asn Glu Thr Met Asn Met Ile Leu Lys Ala  
275 280 285

Lys Leu Gly Met Pro Gln Phe Leu Ser Ala Glu Ala Gln Ser Leu Leu  
290 295 300

Arg Met Leu Phe Lys Arg Asn Pro Ala Asn Arg Leu Gly Ser Glu Gly  
305 310 315 320

Val Glu Glu Ile Lys Arg His Leu Phe Phe Ala Asn Ile Asp Trp Asp  
325 330 335

Lys Leu Tyr Lys Arg Glu Val Gln Pro Pro Phe Lys Pro Ala Ser Gly  
340 345 350

Lys Pro Asp Asp Thr Phe Cys Phe Asp Pro Glu Phe Thr Ala Lys Thr  
355 360 365

Pro Lys Asp Ser Pro Gly Leu Pro Ala Ser Ala Asn Ala His Gln Leu  
370 375 380

Phe Lys Gly Phe Ser Phe Val Ala Thr Ser Ile Ala Glu Glu Tyr Lys  
385 390 395 400

Ile Thr Pro Ile Thr Ser Ala Asn Val Leu Pro Ile Val Gln Ile Asn  
405 410 415

Gly Asn Ala Ala Gln Phe Gly Glu Val Tyr Glu Leu Lys Glu Asp Ile  
420 425 430

Gly Val Gly Ser Tyr Ser Val Cys Lys Arg Cys Ile His Ala Thr Thr  
435 440 445

Asn Met Glu Phe Ala Val Lys Ile Ile Asp Lys Ser Lys Arg Asp Pro  
450 455 460

Ser Glu Glu Ile Glu Ile Leu Met Arg Tyr Gly Gln His Pro Asn Ile  
465 470 475 480

Ile Thr Leu Lys Asp Val Phe Asp Asp Gly Arg Tyr Val Tyr Leu Val  
485 490 495

Thr Asp Leu Met Lys Gly Gly Glu Leu Leu Asp Arg Ile Leu Lys Gln  
500 505 510

Lys Cys Phe Ser Glu Arg Glu Ala Ser Asp Ile Leu Tyr Val Ile Ser

515		520		525
Lys Thr Val Asp Tyr Leu His Cys Gln Gly Val Val His Arg Asp Leu				
530		535		540
Lys Pro Ser Asn Ile Leu Tyr Met Asp Glu Ser Ala Ser Ala Asp Ser				
545		550		555
				560
Ile Arg Ile Cys Asp Phe Gly Phe Ala Lys Gln Leu Arg Gly Glu Asn				
		565		570
				575
Gly Leu Leu Leu Thr Pro Cys Tyr Thr Ala Asn Phe Val Ala Pro Glu				
		580		585
				590
Val Leu Met Gln Gln Gly Tyr Asp Ala Ala Cys Asp Ile Trp Ser Leu				
		595		600
				605
Gly Val Leu Phe Tyr Thr Met Leu Ala Gly Tyr Thr Pro Phe Ala Asn				
		610		615
				620
Gly Pro Asn Asp Thr Pro Glu Glu Ile Leu Leu Arg Ile Gly Asn Gly				
		625		630
				635
Lys Phe Ser Leu Ser Gly Gly Asn Trp Asp Asn Ile Ser Asp Gly Ala				
		645		650
				655
Lys Asp Leu Leu Ser His Met Leu His Met Asp Pro His Gln Arg Tyr				
		660		665
				670
Thr Ala Glu Gln Ile Leu Lys His Ser Trp Ile Thr His Arg Asp Gln				
		675		680
				685
Leu Pro Asn Asp Gln Pro Lys Arg Asn Asp Val Ser His Val Val Lys				
		690		695
				700
Gly Ala Met Val Ala Thr Tyr Ser Ala Leu Thr His Lys Thr Phe Gln				
		705		710
				715
Pro Val Leu Glu Pro Val Ala Ala Ser Ser Leu Ala Gln Arg Arg Ser				
		725		730
				735
Met Lys Lys Arg Thr Ser Thr Gly Leu				
		740		745

<210> 5  
<211> 22

<212> DNA  
<213> Homo sapiens

<400> 5  
ggagcaaagg gagcaatggt tg 22

<210> 6  
<211> 22  
<212> DNA  
<213> Homo sapiens

<400> 6  
tcttcatcca gtttggccta gg 22

<210> 7  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 7  
tgggtgtgaa ccatgagaag 20

<210> 8  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 8  
gtgtcgctgt tgaagtcaga 20